



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

$OP$ ,  $OQ$ , and  $OR$ , we have

$$OP^2(\cos 2\alpha + i \sin 2\alpha) + OQ^2(\cos 2\alpha' + i \sin 2\alpha') = -OR^2(\cos 2\alpha'' + i \sin 2\alpha''),$$

from datum; hence,

$$OP^2 \cos 2\alpha + OQ^2 \cos 2\alpha' = -OR^2 \cos 2\alpha'' \quad (1)$$

and

$$OP^2 \sin 2\alpha + OQ^2 \sin 2\alpha' = -OR^2 \sin 2\alpha''. \quad (2)$$

Now  $P$  and  $Q$  being points on the ellipse, we have from known properties,

$$OP^2 \cos^2 \alpha + OQ^2 \cos^2 \alpha' = a^2, \quad OP^2 \sin^2 \alpha + OQ^2 \sin^2 \alpha' = b^2;$$

hence,  $OP^2 \cos 2\alpha + OQ^2 \cos 2\alpha' = a^2 - b^2$ , and (1) becomes

$$-OR^2 \cos 2\alpha'' = a^2 - b^2. \quad (3)$$

Also, from known properties concerning the ends of conjugate diameters,

$$OP^2 \sin 2\alpha = -OQ^2 \sin 2\alpha';$$

hence, (2) becomes

$$-OR^2 \sin 2\alpha'' = 0. \quad (4)$$

It follows from (3) and (4), that  $2\alpha'' = 180^\circ$  or  $540^\circ$ , and  $OR^2 = a^2 - b^2$ , that is,  $OR = \sqrt{a^2 - b^2}$ , the distance from the center to focus, and  $\alpha'' = 90^\circ$  or  $270^\circ$ , which shows that  $OR$  lies on the minor axis.

Also solved by H. HALPERIN, A. M. HARDING, and H. L. OLSON.

**2780 [1919, 311].** Proposed by ELMER LATSHAW, West Philadelphia, Pa.

A quadrilateral whose sides are  $a, 2a, 3a, 4a$  is inscribed in a circle. Find the radius of the circle.

I. SOLUTION BY H. S. UHLER, Yale University.

The interest in this problem may be enhanced by giving a perfectly general solution. Let the sides of any convex inscriptible quadrilateral be denoted by  $a_1, a_2, a_3, a_4$ . A diagonal  $c$  may be drawn dividing the quadrilateral into two non-overlapping triangles the sides of which are  $a_1, a_2, c$  and  $a_3, a_4, c$ , respectively. If the angle between  $a_1$  and  $a_2$  be symbolized by  $C$ , the angle between  $a_3$  and  $a_4$  must be  $180^\circ - C$ . Accordingly

$$c^2 = a_1^2 + a_2^2 - 2a_1a_2 \cos C,$$

$$c^2 = a_3^2 + a_4^2 + 2a_3a_4 \cos C.$$

Eliminating  $2\cos C$  we find

$$c^2 = \frac{(a_1a_3 + a_2a_4)(a_2a_3 + a_4a_1)}{a_1a_2 + a_3a_4}. \quad (1)$$

The area of a plane triangle having the sides  $a_1, a_2, c$  is given by either member of the following equation

$$\frac{a_1a_2c}{4R} = \sqrt{s(s-a_1)(s-a_2)(s-c)}, \quad (2)$$

where  $2s = a_1 + a_2 + c$ , and  $R$  denotes the radius of the circumscribed circle.

Substituting the trinomial value of  $s$  in equation (2) we obtain

$$\frac{a_1a_2c}{R} = \sqrt{[(a_1 + a_2)^2 - c^2][c^2 - (a_1 - a_2)^2]}. \quad (3)$$

Replacing  $c$  in equation (3) by expression (1) we eventually find that

$$R = \frac{\sqrt{(a_1a_2 + a_3a_4)(a_1a_3 + a_4a_2)(a_2a_3 + a_4a_1)}}{\sqrt{(a_2 + a_3 + a_4 - a_1)(a_3 + a_4 + a_1 - a_2)(a_4 + a_1 + a_2 - a_3)(a_1 + a_2 + a_3 - a_4)}}, \quad (4)$$

or

$$R = \frac{1}{4K} \sqrt{(a_1a_2 + a_3a_4)(a_1a_3 + a_4a_2)(a_2a_3 + a_4a_1)}. \quad (5)$$

where if  $2S = a_1 + a_2 + a_3 + a_4$ ,  $K = \sqrt{(S - a_1)(S - a_2)(S - a_3)(S - a_4)} = \text{area of quadrilateral.}$

The denominator of formula (4) brings out the geometrically-evident fact that each side of the quadrilateral must not exceed the sum of the remaining three sides. When the quadrilateral degenerates into a straight line, formulas (4) and (5) give  $R = \infty$ , as they should. These formulæ also show explicitly that the order or succession of the sides has no effect on the value of  $R$ , a fact which is obvious geometrically since the sum of the arcs subtended by the four sides of the quadrilateral equals the entire circumference.

The answer to the given problem may be obtained at once from formula (5) by substituting  $a, 2a, 3a, 4a, 5a$  for  $a_1, a_2, a_3, a_4, S$  respectively. It is

$$R = \frac{a\sqrt{385}}{4\sqrt{6}} = (2.002602\cdots)a.$$

## II. SOLUTION BY BING CHIN WONG, Berkeley, Calif.

Let  $ABCD$  be the polygon with sides  $AB = a, BC = 2a, CD = 3a, DA = 4a$  inscribed in the circle with  $O$  as center. Join  $O$  to  $A, B, C, D$ . Then

$$\angle AOB + \angle BOC + \angle COD + \angle DOA = 2\alpha + 2\beta + 2\gamma + 2\delta = 2\pi,$$

or

$$\alpha + \beta + \gamma + \delta = \pi.$$

Then

$$\cos(\alpha + \beta) + \cos(\gamma + \delta) = 0,$$

or

$$\cos \alpha \cos \beta - \sin \alpha \sin \beta + \cos \gamma \cos \delta - \sin \gamma \sin \delta = 0. \quad (\text{I})$$

Let  $r$  be the radius of the circle. We obtain from the figure,

$$\sin \alpha = \frac{a}{2r}, \quad \sin \beta = \frac{a}{r}, \quad \sin \gamma = \frac{3a}{2r}, \quad \sin \delta = \frac{2a}{r};$$

and

$$\cos \alpha = \frac{\sqrt{4r^2 - a^2}}{2r}, \quad \cos \beta = \frac{\sqrt{r^2 - a^2}}{r}, \quad \cos \gamma = \frac{\sqrt{4r^2 - 9a^2}}{2r}, \quad \cos \delta = \frac{\sqrt{r^2 - 4a^2}}{r}.$$

Substituting these values in (I) and multiplying by  $2r^2$ , we have

$$\sqrt{4r^2 - a^2}\sqrt{r^2 - a^2} + \sqrt{4r^2 - 9a^2}\sqrt{r^2 - 4a^2} = 7a^2.$$

Squaring, collecting terms, and dividing by 2, we have

$$\sqrt{(4r^2 - a^2)(r^2 - a^2)(r^2 - 4a^2)(4r^2 - 9a^2)} = 6a^4 + 15a^2r^2 - 4r^4.$$

Squaring again and collecting terms, we have

$$96a^4r^4 = 385a^6r^2, \quad \text{or} \quad r^2 = 385a^2/96,$$

and, therefore,

$$r = a\sqrt{385/96} = a\sqrt{2310/24} = 2.0026a.$$

Also solved by H. C. BRADLEY, H. N. CARLETON, S. A. COREY, LAURA GUGGENBUHL, T. F. NOISMANN, H. L. OLSON, A. PELLETIER, J. B. REYNOLDS, and the Proposer.

**2781 [1919, 311]. Proposed by J. L. RILEY, Stephenville, Texas.**

Show that the asymptotic lines on a pseudospherical surface are curves of constant torsion.<sup>1</sup>

**SOLUTION BY OTTO DUNKEL, Washington University.**

The Theorem of Eüneper states that the square of the torsion of an asymptotic line at any point of a surface is equal to the total curvature of the surface (with sign changed) at the point (Eisenhart, *Differential Geometry*, p. 140). Since it is known that the total curvature of a pseudospherical surface is constant, it follows at once that the torsion is constant.

<sup>1</sup> This problem is given as an example in Eisenhart's *Differential Geometry*, p. 290.—EDITORS.